arXiv:1612.01698v1 [math.NT] 6 Dec 2016

ON CERTAIN MOMENTS OF HARDY'S FUNCTION $Z(t)$ OVER SHORT INTERVALS

Aleksandar Ivić

Abstract. Let as usual $Z(t) = \zeta(\frac12 + it)\chi^{-1/2}(\frac12 + it)$ denote Hardy's function, where $\zeta(s) = \chi(s)\zeta(1-s)$. Assuming the Riemann hypothesis upper and lower bounds for some integrals involving $Z(t)$ and $Z'(t)$ are proved. It is also proved that

$$H(\log T)^{k^2} \ll_{k,\alpha} \sum_{T<\gamma\leqslant T+H} \max_{\gamma\leqslant\tau_\gamma\leqslant\gamma^+} |\zeta(\tfrac12 + i\tau_\gamma)|^{2k} \ll_{k,\alpha} H(\log T)^{k^2}.$$

Here $k > 1$ is a fixed integer, γ, γ^+ denote ordinates of consecutive complex zeros of $\zeta(s)$ and $T^\alpha \leqslant H \leqslant T$, where α is a fixed constant such that $0 < \alpha \leqslant 1$. This sharpens and generalizes a result of M.B. Milinovich [17].

1. Introduction

Let the Riemann zeta-function be, as usual,

$$\zeta(s) = \sum_{n=1}^\infty n^{-s} \qquad (\Re s > 1).$$

For $\Re s \leqslant 1$ one defines $\zeta(s)$ by analytic continuation (see the monographs of H.M. Edwards [3], the author [11] and E.C. Titchmarsh [20] for the properties of $\zeta(s)$). Here the Riemann Hypothesis (RH), that all complex zeros of $\zeta(s)$ satisfy $\Re s = \frac12$, is assumed *throughout the paper*. The Riemann zeta-function satisfies the functional equation

$$\zeta(s) = \chi(s)\zeta(1-s) \quad (\forall s) \in \mathbb{C}), \quad \chi(s) := \frac{\Gamma(\frac12(1-s))}{\Gamma(\frac12 s)}\pi^{s-1/2}, \tag{1.1}$$

1991 *Mathematics Subject Classification.* 11M06.

Key words and phrases. Riemann zeta-function, Riemann Hypothesis, Hardy's function, moments, short intervals.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$-TEX

where $\Gamma(s)$ is the familiar gamma-function. One then defines Hardy's function $Z(t)$ as

$$Z(t) := \zeta(\tfrac12 + it)\bigl(\chi(\tfrac12 + it)\bigr)^{-1/2}, \tag{1.2}$$

which is real for t real and $|\zeta(\frac12 + it)| = |Z(t)|$. Thus the real zeros of $Z(t)$ correspond to the zeros of $\zeta(s)$ of the form $\frac12 + it$, which makes Hardy's function an invaluable tool in the study of zeros of $\zeta(s)$ on the critical line $\Re s = \frac12$. For an extensive account on $Z(t)$ the reader is referred to the author's monograph [16].

Several papers deal with the estimation of the sum

$$\mathcal{M}_k(T) := \frac{1}{N(T)} \sum_{0<\gamma\leqslant T} \max_{\gamma\leqslant\tau_\gamma\leqslant\gamma^+} |\zeta(\tfrac12 + i\tau_\gamma)|^{2k} \equiv \frac{1}{N(T)} \sum_{0<\gamma\leqslant T} \max_{\gamma\leqslant\tau_\gamma\leqslant\gamma^+} |Z(\tau_\gamma)|^{2k}. \tag{1.3}$$

Here $k \in \mathbb{N}$ is fixed, and γ, γ^+ denote ordinates of consecutive complex zeros of $\zeta(s)$, ordered according to their size. Also, as usual,

$$N(T) = \sum_{0<\gamma\leqslant T} 1 = \frac{T}{2\pi}\log\frac{T}{2\pi} - \frac{T}{2\pi} + O(\log T) \tag{1.4}$$

counts (with multiplicities) the number of zeros of $\zeta(s)$ whose ordinates γ satisfy $0 < \gamma \leqslant T$.

In [2] B. Conrey and A. Ghosh proved, under the RH,

$$\mathcal{M}_1(T) = \frac{\mathrm{e}^2 - 5}{2}\log T + O(1). \tag{1.5}$$

Actually, they prove a somewhat stronger result than (1.5), namely

$$\sum_{T<\gamma\leqslant T+H} \max_{\gamma\leqslant\tau_\gamma\leqslant\gamma^+} |\zeta(\tfrac12 + i\tau_\gamma)|^2 = \frac{\mathrm{e}^2 - 5}{4\pi} H\log^2 T + O(H\log T) \tag{1.6}$$

with $H = T^{3/4}$. This follows from their proof on noting that (1.4) implies

$$N(T+H) - N(T) \ \sim\ \frac{H}{2\pi}\log T \qquad (T\to\infty).$$

B. Conrey [1] obtained, also under the RH,

$$\frac{\sqrt{21}}{45\pi}(1+o(1))\log^4 T \leqslant \mathcal{M}_2(T) \leqslant \frac{1+o(1)}{\pi\sqrt{15}}\log^4 T \quad (T\to\infty), \tag{1.7}$$

and R.R. Hall [4], [5] obtained some further improvements of (1.7). A general result, due to M.B. Milinovich [17], states that under the RH, for fixed $k \in \mathbb{N}$.

$$(\log T)^{k^2-\varepsilon} \ll_{k,\varepsilon} \mathcal{M}_k(T) \ll_{k,\varepsilon} (\log T)^{k^2+\varepsilon}. \tag{1.8}$$

Here $\ll_{k,\varepsilon}$ means that the constant implied by the $\ll$-symbol depends only on k and ε, an arbitrarily small positive number, not necessarily the same one at each occurrence. The bounds in (1.8), when $k = 1, 2$, are implied by (1.6) and (1.7), respectively.

Acknowledgment. The author wishes to thank M.B. Milinovich for valuable remarks.

2. Statement of results

Milinovich [17] derives (1.8) from upper and lower bounds involving certain integrals with $Z(t)$ and $Z'(t)$, which seem to be of independent interest. He investigated integrals over the "long" interval $[0, T]$, but here we are interested in the integrals over the "short" intervals $[T, T+H]$, where $H = H(T)$ may be much smaller than T. We shall prove here the following theorems.

THEOREM 1. *Let $k \geqslant 2$ be a fixed integer. Under the RH we have, for $T^\alpha \leqslant H = H(T) \leqslant T, 0 < \alpha \leqslant 1$ a fixed constant,*

$$\int_T^{T+H} (Z'(t))^2 Z^{2k-2}(t)\,\mathrm{d}t \gg_{k,\alpha} H(\log T)^{k^2+2}. \tag{2.1}$$

THEOREM 2. *Let $k \in \mathbb{N}$ be fixed. Under the RH we have, for $T^\alpha \leqslant H = H(T) \leqslant T, 0 < \alpha \leqslant 1$ a fixed constant,*

$$\int_T^{T+H} |\zeta'(\tfrac{1}{2}+it)|^{2k}\,\mathrm{d}t \ll_{k,\alpha} H(\log T)^{k^2+2}, \tag{2.2}$$

and

$$\int_T^{T+H} \big(Z'(t)\big)^{2k}\,\mathrm{d}t \ll_{k,\alpha} H(\log T)^{k^2+2}. \tag{2.3}$$

These bounds differ from the analogous results of [17] in two aspects. Firstly, Milinovich has the integrals over $[0, T]$, which corresponds to the case $H = T$ in our theorems. Indeed, if (2.1)–(2.3) hold with $H = T$, then replacing T by $T/2, T/2^2, \ldots$ etc. and adding up all the results we obtain (2.1)–(2.3) with the interval of integration $[0, T]$. Secondly, in (2.1) Milinovich obtained $k^2 + 2 - \varepsilon$

as the exponent of the logarithm, and in (2.2) and (2.3) he had the exponents $k^2+2+\varepsilon$ of the logarithm. He remarks on page 1122 that one can get rid of the ε's in his bounds provided that one has

$$\int_0^T |\zeta(\tfrac12+it)|^{2k}\,\mathrm{d}t \ \ll_k \ T(\log T)^{k^2} \tag{2.4}$$

and

$$\int_0^T |\zeta'(\tfrac12+it)|^{2k}\,\mathrm{d}t \ \ll_k \ T(\log T)^{k^2+2k}. \tag{2.5}$$

The estimates (2.4) and (2.5) do hold indeed. Namely K. Soundararajan [19] proved (2.4) with the exponent of $\log T$ in (2.4) equal to $k^2+\varepsilon$. The author [15] improved and sharpened Soundararajan's bound by showing that

$$\int_T^{T+H} |\zeta(\tfrac12+it)|^{2k}\,\mathrm{d}t \ \ll_{k,\alpha} \ H(\log T)^{k^2\left(1+O(1/\log_3 T)\right)} \quad \text{(RH)}. \tag{2.6}$$

Here $T^\alpha \leqslant H \leqslant T$ where $0<\alpha\leqslant 1$ is a fixed number, and

$$\log_3 T = \log\log\log T = \log(\log_2 T).$$

Note that [15] appeared before [19] because of the backlog of "Ann. Math." The key result in [15], which is a proper generalization of the corresponding result in [19], is

THEOREM A. *Let $H=T^\theta$ where $0<\theta\leqslant 1$ is a fixed number, and let $\mu(T,H,V)$ denote the measure of points t from $[T-H,\,T+H]$ such that*

$$\log|\zeta(\tfrac12+it)| \geqslant V, \quad 10\sqrt{\log_2 T} \leqslant V \leqslant \frac{3\log 2T}{8\log_2(2T)}.$$

Then, under the RH, for $10\sqrt{\log_2 T}\leqslant V\leqslant \log_2 T$ we have

$$\mu(T,H,V) \ \ll \ H\frac{V}{\sqrt{\log_2 T}}\exp\left(-\frac{V^2}{\log_2 T}\Big(1-\frac{7}{2\theta\log_3 T}\Big)\right),$$

for $\log_2 T\leqslant V\leqslant \frac12\theta\log_2 T\log_3 T$ we have

$$\mu(T,H,V) \ \ll \ H\exp\left(-\frac{V^2}{\log_2 T}\Big(1-\frac{7V}{4\theta\log_2 T\log_3 T}\Big)^2\right),$$

and for $\frac{1}{2}\theta\log_2 T\log_3 T \leqslant V \leqslant \frac{3\log 2T}{8\log_2(2T)}$ *we have*

$$\mu(T,H,V) \ll H\exp(-\tfrac{1}{20}\theta V\log V).$$

Later A. Harper [9] (RH) improved the upper bound in [19] by establishing (2.4) for the long interval $[0,T]$. As remarked in [9] on p. 4, the method of [15] leading to (2.6), i.e., Theorem A, can be combined with that of [9] to produce the sharp upper bound over the short interval $[T,T+H]$, namely

$$\int_T^{T+H} |\zeta(\tfrac{1}{2}+it)|^{2k}\,\mathrm{d}t \ll_{k,\alpha} H(\log T)^{k^2} \quad \text{(RH)}. \tag{2.7}$$

The bound in (2.7) is the key ingredient in the proof of our results. It is, up to the values of the $\ll$-constants, best possible, since long ago it was shown by R. Balasubramanian and K. Ramachandra (see the latter's monograph [18], in particular the remark on p. 45) that, if $k \geqslant 1$ is a fixed integer, then for $C(\varepsilon,k)\log\log T \leqslant H \leqslant T/2$ we have

$$\int_T^{T+H} |\zeta(\tfrac{1}{2}+it)|^{2k}\,\mathrm{d}t \geqslant (C_k'-\varepsilon)H(\log H)^{k^2}, \tag{2.8}$$

where

$$C_k' = \frac{1}{2\Gamma(k^2+1)}\prod_p\Big\{(1-p^{-1})^{k^2}\sum_{m=0}^{\infty}\Big(\frac{\Gamma(k+m)}{\Gamma(k)m!}\Big)^2 p^{-m}\Big\}.$$

We note that the lower bound in (2.8) is *unconditional*, with a very wide range for H. As for (2.5), it will be shown later that a corresponding result holds over $[T,T+H]$.

THEOREM 3. *Let* $1 < k \in \mathbb{N}$ *be fixed,* γ,γ^+ *denote ordinates of consecutive complex zeros of* $\zeta(s)$ *and* $T^\alpha \leqslant H = H(T) \leqslant T$*, where* α *is a fixed constant such that* $0 < \alpha \leqslant 1$*. Under the RH we have then*

$$H(\log T)^{k^2} \ll_{k,\alpha} \sum_{T<\gamma\leqslant T+H}\ \max_{\gamma\leqslant\tau_\gamma\leqslant\gamma^+} |\zeta(\tfrac{1}{2}+i\tau_\gamma)|^{2k} \ll_{k,\alpha} H(\log T)^{k^2}. \tag{2.9}$$

Remark. The case $k=1$ was treated in [2] (see (1.6)) and is not covered by Theorem 3. This is because Theorem 1 does not cover the case $k=1$. The method of its proof (see (3.4)) does not work in obtaining a lower bound for

$$\int_T^{T+H} |Z'(t)Z(t)|\,\mathrm{d}t.$$

3. Proof of Theorem 1

As was also done in [Mil], we follow Conrey and Ghosh [2], and introduce the analytic function

$$Z_1(s) := \zeta'(s) - \frac{\chi'(s)}{2\chi(s)}\zeta(s). \tag{3.1}$$

Its usefulness comes from the fact that differentiation of (1.2) gives

$$Z'(t) = i\left\{\zeta'(\tfrac12+it) - \tfrac12\frac{\chi'(\frac12+it)}{\chi(\frac12+it)}\zeta(\tfrac12+it)\right\}\chi^{-1/2}(\tfrac12+it). \tag{3.2}$$

This gives

$$|Z'(t)| \ = \ |Z_1(\tfrac12+it)|. \tag{3.3}$$

Using (3.3) and $|Z(t)| = |\zeta(\frac12+it)|$ one may write

$$\int_T^{T+H}(Z'(t))^2Z^{2k-2}(t)\,\mathrm{d}t = \int_T^{T+H}\left|Z_1(\tfrac12+it)\zeta(\tfrac12+it)^{k-1}\right|^2\mathrm{d}t. \tag{3.4}$$

The basic idea is to use the inequality

$$\begin{aligned}&\Big|\int_T^{T+H}Z_1(\tfrac12+it)\zeta(\tfrac12+it)^{k-1}\bar A(t)\,\mathrm{d}t\Big|^2\\ &\leqslant \int_T^{T+H}(Z'(t))^2Z^{2k-2}(t)\,\mathrm{d}t\cdot\int_T^{T+H}|A(t)|^2\,\mathrm{d}t.\end{aligned} \tag{3.5}$$

This comes on using (3.4) and the Cauchy-Schwarz inequality for integrals with a suitably chosen function $A(t)$. Following [Mil] we set

$$A(t) := \mathcal{A}(\tfrac12+it),\quad \mathcal{A}(s) = \mathcal{A}(s;k,\xi) := \sum_{n\leqslant\xi}d_k(n)n^{-s},$$

where $d_k(n)$ (generated by $\zeta^k(s)$ for $\Re s>1$) is the (generalized) divisor function which represents the number of ways n can be written as a product of k fixed factors (see e.g., Chapter 13 of [11] for more properties). The parameter ξ is given by $\xi = T^\theta, 0<\theta<1$. The function $A(t)$ has the property that in mean square it behaves like $(\log\xi)^{k^2}$ (see Chapter 13 of [11]). Therefore, by the mean value theorem for Dirichlet polynomials (see e.g., Theorem 5.2 of [11]), we have

$$\begin{aligned}\int_T^{T+H}|A(t)|^2\,\mathrm{d}t &= H\sum_{n\leqslant\xi}d_k^2(n)n^{-1} + O\Big(\sum_{n\leqslant\xi}d_k^2(n)\Big)\\ &= H(C_k+o(1))(\log\xi)^{k^2} + O(\xi(\log\xi)^{k^2-1})\end{aligned} \tag{3.6}$$

for $2 \leqslant \xi \leqslant T$ and a positive constant C_k, which may be made explicit. It remains to estimate from below

$$\begin{aligned}&\int_T^{T+H} Z_1(\tfrac{1}{2}+it)\zeta(\tfrac{1}{2}+it)^{k-1}\bar{A}(t)\,\mathrm{d}t\\ &= \frac{1}{i}\int_{1/2+iT}^{1/2+iT+iH} Z_1(s)\zeta^k(s)\mathcal{A}(1-s)\,\mathrm{d}s\\ &= \int_{a+iT}^{a+iT+iH} \frac{1}{i}Z_1(s)\zeta^k(s)\mathcal{A}(1-s)\,\mathrm{d}s + O_\varepsilon(T^\varepsilon\xi).\end{aligned} \tag{3.7}$$

Here we used Cauchy's theorem and set $a := 1 + 1/\log T$. We also used standard consequences of the RH (see Chapter 12 of [20]):

$$\zeta(s) \ll_{\varepsilon,\sigma} |t|^\varepsilon, \quad \zeta'(s) \ll_{\varepsilon,\sigma} |t|^\varepsilon \qquad (s = \sigma + it, \sigma \geqslant \tfrac{1}{2}),$$

as well as the unconditional, elementary bound $d_k(n) \ll_{\varepsilon,k} n^\varepsilon$ and

$$\frac{\chi'(\sigma+it)}{\chi(\sigma+it)} = -\log\frac{t}{2\pi} + O\Big(\frac{1}{t}\Big). \tag{3.8}$$

One obtains (3.8) by logarithmic differentiation of (1.1) and the use of Stirling's formula for the gamma-function. It is valid for $\frac{1}{2} \leqslant \Re s \leqslant 2$, and the O-term in (3.8) in fact admits a full asymptotic expansion in terms of negative exponents of t, and the left-hand side of (3.8) can be further differentiated. The integral on the right-hand side of (3.7) is written as $J_1 + J_2$, where

$$\begin{aligned}J_1 :=&\ \frac{1}{i}\int_{a+iT}^{a+iT+iH} \zeta'(s)\zeta^{k-1}\mathcal{A}(1-s)\,\mathrm{d}s,\\ J_2 :=&\ -\frac{1}{2i}\int_{a+iT}^{a+iT+iH} \frac{\chi'(s)}{\chi(s)}\zeta^k(s)\mathcal{A}(1-s)\,\mathrm{d}s.\end{aligned}$$

Similarly as in [17] one shows that

$$J_1 = -H\sum_{n\leqslant\xi} \tilde{d}_k(n)d_k(n)n^{-1} + O(T^\varepsilon\xi)$$

with

$$\tilde{d}_k(n) := \sum_{\delta|n} d_{k-1}(\delta)\log\frac{n}{\delta} \leqslant d_k(n)\log n,$$

and

$$J_2 = \tfrac{1}{2}H\left(\log\frac{T}{2\pi\mathrm{e}}\right)\sum_{n\leqslant\xi} d_k^2(n)n^{-1} + O(T^\varepsilon\xi).$$

This yields

$$\begin{aligned}
&\int_T^{T+H} Z_1(\tfrac12+it)\zeta(\tfrac12+it)^{k-1}\bar A(t)\,\mathrm{d}t = J_1+J_2+O(T^\varepsilon\xi)\\
&= -H\sum_{n\leqslant\xi}\tilde d_k(n)d_k(n)n^{-1}+O(T^\varepsilon\xi)+\tfrac12 H\log\frac{T}{2\pi}(1+o(1))\sum_{n\leqslant\xi}d_k^2(n)n^{-1}\\
&\geqslant -H\sum_{n\leqslant\xi}d_k^2(n)n^{-1}\log n+\tfrac12 H\log\frac{T}{2\pi}(1+o(1))\sum_{n\leqslant\xi}d_k^2(n)n^{-1}+O(T^\varepsilon\xi)\\
&\geqslant \Big\{\tfrac12 H(1+o(1))\log\frac{T}{2\pi}-H\log\xi\Big\}\sum_{n\leqslant\xi}d_k^2(n)n^{-1}+O(T^\varepsilon\xi)\\
&\geqslant A_k H\log T\cdot(\log\xi)^{k^2}
\end{aligned}$$

for $\xi=T^\theta,\theta=\frac12\alpha$ and ε sufficiently small. From (3.5) and (3.6) we finally gather that

$$H^2\log^2 T(\log T)^{2k^2}\ll_{k,\alpha}\int_T^{T+H}(Z'(t))^2Z^{2k-2}(t)\,\mathrm{d}t\cdot H(\log T)^{k^2},$$

and (2.1) of Theorem 1 follows.

4. Proof of Theorem 2

First note that, for $0<R\leqslant\frac12, T\leqslant t\leqslant 2T$, by Cauchy's integral formula we have

$$\zeta'(\tfrac12+it)\;=\;\frac{1}{2\pi i}\int_{|z|=R}\frac{\zeta(\frac12+it+z)}{z^2}\,\mathrm{d}z.$$

This yields

$$\int_T^{T+H}|\zeta'(\tfrac12+it)|^{2k}\,\mathrm{d}t=\frac{1}{(2\pi)^{2k}}\int_T^{T+H}\left|\int_{|z|=R}\frac{\zeta(\frac12+it+z)}{z^2}\,\mathrm{d}z\right|^{2k}\mathrm{d}t. \tag{4.1}$$

By Hölder's inequality for integrals the right-hand side of (4.1) does not exceed

$$\begin{aligned}
&\frac{1}{(2\pi)^{2k}}\int_T^{T+H}\left\{\int_{|z|=R}|\zeta(\tfrac12+it+z)|^{2k}|\,\mathrm{d}z|\right\}\cdot\left\{\frac{|\,\mathrm{d}z|}{|z|^{4k/(2k-1)}}\right\}^{2k-1}\mathrm{d}t\\
&\leqslant\frac{1}{(2\pi)^{2k}}\int_T^{T+H}\left\{\int_{|z|=R}|\zeta(\tfrac12+it+z)|^{2k}|\,\mathrm{d}z|\right\}(2\pi R)^{2k-1}R^{-4k}\\
&\leqslant\frac{1}{R^{2k}}\max_{0\leqslant\theta\leqslant2\pi}\int_T^{T+H}|\zeta(\tfrac12+it+R\mathrm{e}^{i\theta})|^{2k}\,\mathrm{d}t.
\end{aligned}$$

Therefore

$$\int_T^{T+H} |\zeta'(\tfrac{1}{2}+it)|^{2k}\,\mathrm{d}t \leqslant \frac{1}{R^{2k}} \max_{0\leqslant\theta\leqslant 2\pi} \int_T^{T+H} |\zeta(\tfrac{1}{2}+it+R\mathrm{e}^{i\theta})|^{2k}\,\mathrm{d}t. \tag{4.2}$$

As in [Mil], we could have obtained an inequality for the ℓ-th derivative of $\zeta(\frac{1}{2}+it)$, but this is not necessary for our purposes.

Henceforth let $R = 1/\log T$ in (4.2). The integral on the right-hand side of (4.2) equals

$$\int_T^{T+H} |\zeta\big(\tfrac{1}{2}+R\cos\alpha+i(t+R\sin\alpha)\big)|^{2k}\,\mathrm{d}t. \tag{4.3}$$

Recall that, under the RH (see [Tit]),

$$\zeta(\sigma+it) \ll \exp\left(C\frac{\log t}{\log\log t}\right) \qquad (\tfrac{1}{2}\leqslant\sigma\leqslant 1, C>0, |t|\geqslant 2). \tag{4.4}$$

When $\cos\alpha \geqslant 0$ in (4.3), we use (4.4) to obtain that the integral in (4.3) is equal to

$$\int_T^{T+H} |\zeta(\tfrac{1}{2}+R\cos\alpha+it)|^{2k}\,\mathrm{d}t + o(T). \tag{4.5}$$

When $\pi/2 \leqslant \theta \leqslant 3\pi/2$ in (4.5), that is, when $\cos\alpha \leqslant 0$, we use the functional equation (1.1). In this case we have, with $\sigma = \Re s = \frac{1}{2} + R\cos\alpha, R = 1/\log T$,

$$\chi(s) \ll |t|^{1/2-\sigma} \ll T^{-\cos\alpha/\log T} \ll 1,$$

thus we reduce the estimation of our integral to the case when $\cos\alpha \geqslant 0$. For this we shall use a convexity result which shows that essentially the integral in question is bounded by the $2k$-th moment of $|\zeta(\frac{1}{2}+it)|$ over a short interval. More precisely, let $1/2 \leqslant \sigma \leqslant 3/4, k>0, t\geqslant 2$,

$$J_k(\sigma) := \int_{-\infty}^{\infty} |\zeta(\sigma+it)|^{2k} w_k(t)\,\mathrm{d}t, \quad w_k(t) := \int_T^{T+H} \mathrm{e}^{-2k(t-\tau)^2}\,\mathrm{d}\tau. \tag{4.6}$$

Then

$$J_k(\sigma) \ll T^{\sigma-1/2}\Big(J_k(\tfrac{1}{2})\Big)^{3/2-\sigma} + \mathrm{e}^{-kT^2/4}. \tag{4.7}$$

The bound in (4.7) is the analogue of Lemma 4.2 of [17] for short intervals. This in turn is a result of D.R. Heath-Brown [10], which is also expounded in [12], pp.

321-323. In the original version the interval of integration in the kernel function function $w_k(t)$ was $[T, 2T]$. However, the change made in (4.6) does not affect the proof, and one obtains (4.7). Now note that $w_k(t) \gg 1$ for $t \in [T, T+H]$, so that

$$\int_T^{T+H} |\zeta(\sigma+it)|^{2k}\,\mathrm{d}t \ll J_k(\sigma). \tag{4.8}$$

We have the bound $w_k(t) \ll \exp(-2kH^2)$ when $t \leqslant T-H$ or $t \geqslant T+2H$. On the other hand $w_k(t) \ll \exp(-kt^2)$ for $t<0$ or $t>3T$. Thus combining (4.7) and (4.8) it follows that

$$\begin{aligned}\int_T^{T+H} |\zeta(\sigma+it)|^{2k}\,\mathrm{d}t &= \int_{-\infty}^0 + \int_0^{T-H} + \int_{T-H}^{T+2H} + \int_{T+2H}^{\infty}\\ &\ll 1 + \int_{T-H}^{T+2H} |\zeta(\tfrac12+it)|^{2k}\,\mathrm{d}t\\ &\ll_{k,\alpha} H(\log T)^{k^2},\end{aligned} \tag{4.9}$$

where in the last step (2.7) was used. Inserting (4.9) in (4.2) the bound in (2.2) follows. The estimate (2.3) easily follows from (2.2), (2.9) and (3.2). Theorem 2 is proved.

5. Proof of Theorem 3

It is in the folklore that $Z(t)$, for $t \geqslant 14$, cannot have a negative local maximum or a positive local minimum under the RH. For this, see [3], or [13], [14], [6]. In other words, the zeros of $Z(t)$ and $Z'(t)$ are interlacing. Thus if γ, γ^+ are consecutive zeros of $Z(t)$, there is a unique point $\lambda_\gamma \in [\gamma, \gamma^+]$ for which $Z'(\lambda_\gamma)=0$ (this is trivially true if $\gamma=\gamma^+$, that is, if γ is a multiple zero of $Z(t)$). Therefore

$$\max_{\gamma \leqslant \tau_\gamma \leqslant \gamma^+} |\zeta(\tfrac12+i\tau_\gamma)|^{2k} = Z^{2k}(\lambda_\gamma). \tag{5.1}$$

Then, since $Z(t)$ is positive in (γ, λ_γ) and negative in $(\lambda_\gamma, \gamma^+)$, or conversely, we have

$$\begin{aligned}\int_\gamma^{\gamma^+} |Z'(t)Z^{2k-1}(t)|\,\mathrm{d}t &= \left|\int_\gamma^{\lambda_\gamma} Z'(t)Z^{2k-1}(t)\,\mathrm{d}t - \int_{\lambda_\gamma}^{\gamma} Z'(t)Z^{2k-1}(t)\,\mathrm{d}t\right|\\ &= \frac{1}{k}Z^{2k}(\lambda_\gamma).\end{aligned} \tag{5.2}$$

Therefore (5.1) and (5.2) give, in view of (4.4),

$$\sum_{T<\gamma\leqslant T+H} \max_{\gamma\leqslant\tau_\gamma\leqslant\gamma^+} |\zeta(\tfrac12+i\tau_\gamma)|^{2k} = k\int_T^{T+H} |Z'(t)Z^{2k-1}(t)|\,\mathrm{d}t + O_{k,\varepsilon}(T^\varepsilon). \tag{5.3}$$

Assume that $k \geqslant 2$, so that $2k-2 \geqslant 2$. To bound the integral on the right-hand side of (5.3) from below, note that

$$|(Z')^2 Z^{2k-2}| = |Z'|^{1/2}|Z|^{k-1/2} \cdot |Z'|^{3/2} \cdot |Z|^{k-3/2}.$$

Thus Hölder's inequality for integrals shows that

$$\int_T^{T+H} (Z'(t))^2 Z^{2k-2}(t)\,\mathrm{d}t \leqslant$$

$$\left(\int_T^{T+H} (|Z'|^{1/2}|Z|^{k-1/2})^p\,\mathrm{d}t\right)^{\frac{1}{p}} \left(\int_T^{T+H} |Z'|^{3q/2}\,\mathrm{d}t\right)^{\frac{1}{q}} \left(\int_T^{T+H} |Z|^{r(k-3/2)}\,\mathrm{d}t\right)^{\frac{1}{r}}$$

with $p, q, r > 0, 1/p + 1/q + 1/r = 1$. Take

$$\frac{1}{p} = \frac{1}{2}, \quad \frac{1}{q} = \frac{3}{4k}, \quad \frac{1}{r} = \frac{1}{2} - \frac{3}{4k}.$$

Then the right-hand side is, on using (2.3) and (2.4),

$$\begin{aligned} &\leqslant \left(\int_T^{T+H} |ZZ|^{2k-1}\,\mathrm{d}t\right)^{\frac{1}{2}} \left(\int_T^{T+H} |Z'|^{2k}\,\mathrm{d}t\right)^{\frac{3}{4k}} \left(\int_T^{T+H} |Z|^{2k}\,\mathrm{d}t\right)^{\frac{1}{2}-\frac{3}{4k}} \\ &\ll_{k,\alpha} \left(\int_T^{T+H} |ZZ|^{2k-1}\,\mathrm{d}t\right)^{\frac{1}{2}} \left(H(\log T)^{k^2+2k}\right)^{\frac{3}{4k}} \left(H(\log T)^{k^2}\right)^{\frac{1}{2}-\frac{3}{4k}}. \end{aligned}$$

This gives, on using (2.1),

$$H(\log T)^{k^2+2} \ll_{k,\alpha} I^{1/2} \left(H(\log T)^{k^2+2k}\right)^{\frac{3}{4k}} \left(H(\log T)^{k^2}\right)^{\frac{1}{2}-\frac{3}{4k}},$$

which on simplifying yields

$$I := \int_T^{T+H} |Z'(t)Z^{2k-1}(t)|\,\mathrm{d}t \gg_{k,\alpha} H(\log T)^{k^2+1}.$$

In view of (5.3) this proves the lower bound in (2.9) of Theorem 3.

As for the upper bound, the integral in (5.3) does not exceed, by Hölder's inequality for integrals,

$$\begin{aligned} &\left|\int_T^{T+H} (Z'(t))^{2k}\,\mathrm{d}t\right|^{1/(2k)} \left|\int_T^{T+H} |\zeta(\tfrac{1}{2}+it)|^{2k}\,\mathrm{d}t\right|^{1-1/(2k)} \\ &\ll_{k,\alpha} \left\{H(\log T)^{k^2+2k}\right\}^{1/(2k)} \left\{H(\log T)^{k^2}\right\}^{1-1/(2k)} \\ &= H(\log T)^{k^2+1}, \end{aligned} \tag{5.4}$$

which finishes the proof of Theorem 3 when $k \geqslant 2$. Here we used (2.3) and (2.4), and we note that the bound in (5.4) holds also for $k = 1$. It is the lower bound in this case which is problematic.

References

[1] J.B. Conrey, The fourth moment of derivatives of the Riemann zeta-function, Quart. J. Math. Oxford Ser. (2)**39**(1988), 21-36.

[2] J.B. Conrey and A. Ghosh, A mean value theorem for the Riemann zeta-function at its relative extrema on the critical line, J. Lond. Math. Soc., II. Ser. **32**(1985), 193-202.

[3] H.M. Edwards, Riemann's zeta-function, Academic Press, New York-London, 1974.

[4] R.R. Hall, The behaviour of the Riemann zeta-function on the critical line, Mathematica **46**(1999), 281-313.

[5] R.R. Hall, On the extreme values of the Riemann zeta-function between its zeros on the critical line, J. reine angew. Math. **560**(2003), 29-41.

[6] R.R. Hall, On the stationary points of Hardy's function $Z(t)$, Acta Arith. **111**(2004), 125-140.

[7] R.R. Hall, A new unconditional result about large spaces between zeta zeros, Mathematika **53**(2005), 101-113.

[8] R.R. Hall, Extreme values of the Riemann zeta-function on short zero intervals, Acta Arith. **121**(2006), 259-273.

[9] A.J. Harper, Sharp conditional bounds for moments of the Riemann zeta-function, preprint available at `arXiv:1305.4618`.

[10] D.R. Heath-Brown, Fractional moments of the Riemann zeta-function, J. London Math. Soc. **24**(1981), 65-78.

[11] A. Ivić, The Riemann zeta-function, John Wiley & Sons, New York 1985 (reissue, Dover, Mineola, New York, 2003).

[12] A. Ivić, Mean values of the Riemann zeta-function, LN's **82**, Tata Inst. of Fundamental Research, Bombay, 1991 (distr. by Springer Verlag, Berlin etc.).

[13] A. Ivić, On some results concerning the Riemann Hypothesis, in "Analytic Number Theory" (Kyoto, 1996) ed. Y. Motohashi, LMS LNS **247**, Cambridge University Press, Cambridge, 1997, pp. 139-167.

[14] A. Ivić, On some reasons for doubting the Riemann Hypothesis, in P. Borwein, S. Choi, B. Rooney and A. Weirathmueller, "The Riemann Hypothesis", CMS Books in Mathematics, Springer, 2008.

[15] A. Ivić, On mean value results for the Riemann Zeta-Function in short intervals, Hardy-Ramanujan J. **32**(2009), 4-23.

[16] A. Ivić, The theory of Hardy's Z-function, Cambridge University Press, Cambridge, 2012, 245pp.

[17] M.B. Milinovich, Moments of the Riemann zeta-function at its relative extrema on the critical line, Bull. London Math. Soc. **43**(2011), 1119-1129.

[18] K. Ramachandra, On the mean-value and omega-theorems for the Riemann zeta-function, LN's **85**, Tata Inst. of Fundamental Research (distr. by Springer Verlag, Berlin etc.), Bombay, 1995.

[19] K. Soundararajan, Moments of the Riemann zeta function, Ann. Math. (2) **170**(2009), 981-993.

[20] E.C. Titchmarsh, The theory of the Riemann zeta-function (2nd edition), Oxford University Press, Oxford, 1986.

Aleksandar Ivić, Katedra Matematike RGF-a Universiteta u Beogradu, Đušina 7, 11000 Beograd, Serbia

E-mail address: `aleksandar.ivic@rgf.bg.ac.rs, aivic_2000@yahoo.com`